

GMP Recombinant Human Interleukin-4

 $\begin{array}{c} 50\mu g: Quantity\\ rHuIL-4-50\mu g: Code\\ G017/LC1/030526/PV: Batch\\ 09/2015: Exp.Date\\ -20^{\circ}C: Storage \end{array}$

CERTIFICATE OF ANALYSIS

| Source | E.coli | | | | |
|-------------------|--|--|--|--|--|
| Appearance | Colourless, clear liquid free of particles | | | | |
| Identity | 1 band at 15kDa as measured by SDS-PAGE/Western Blot | | | | |
| Specific activity | 12.4 x 10 exp6 units / mg compared to NIBSC standard (Bioassay) | | | | |
| Endotoxin content | $< 0.1 \text{EU/}\mu\text{g}$ (LAL) | | | | |
| Protein content | 50±10 μg/vial (Lowry/μBCA) | | | | |
| Trehalose | 6±0.5 mg/ml (HPLC) | | | | |
| Sterility | Absence of growth (FTM (30-35°C)) | | | | |
| | Absence of growth (TSB (20-25°C)) | | | | |
| Abnormal toxicity | No weight loss, no abnormal reaction in mice | | | | |
| General Growth | No weight loss, no abnormal reaction in guinea pigs | | | | |
| Physical state | Freeze-dried | | | | |
| Stability | 12 months at -20°C to -80°C | | | | |
| | At least 3 months after reconstruction when stored at -20°C to -80°C | | | | |
| Reconstruction | Use 500µL water for injection | | | | |
| Packaging unit | 50 μg protein (Lowry test) | | | | |
| Purity | >98% as determined by SDS-PAGE and HPLC | | | | |

GMP certification: GENTAUR rh IL-4 is manufactured in full compliance with cGMP in facilities approved by the Belgian Ministry of Health for the production and storage of medicinal products. The manufacturing process does not involve the use of products of animal origin.

BSE/TSE Declaration: GENTAUR BVBA declares that this recombinant human IL-4 (04-GMPhuIL4) is manufactured under strict GMP controls and certifies that the entire product line is BSE (Bovine Spongiform Encephalopathy) and TSE (Transmissible Spongiform Encephalopathy) free. GENTAUR BVBA manufactures this product in Belgium.

Use GENTAUR rh IL-4 is not an approved medicinal product and ONLY be injected as such to patients after the ethical commission approval of your university.


STABILITY TESTING

NB: Time 0 = Final Container

| | | After 7 days at 37°C | | |
|---------------------------------------|----------------|--|------------------------------|--------|
| TEST | APPLICATION | SPECIFICATION | RESULT | CONCL, |
| Electrophoretical pattern | SDS-PAGE | 1 band between 12 and 16 kDa | 1 band between 12 and 16 kDa | PASS |
| Identity | Western Blot | 1 band between 12 and 16 kDa | 1 band between 12 and 16 kDa | PASS |
| Activity | Bioassay | > 5 10^6 U/mg | 10,9 10^6 U/mg | PASS |
| Abnormal toxicity / General safety | on mice | No weight loss, no abnormal Reaction | Absence of symptoms | PASS |
| Abnormal toxicity / General safety | on guinea-pigs | No weight loss, no abnormal Reaction | Absence of symptoms | PASS |
| Sterility test | FTM 30-35°C | Absence of growth | Absence of growth | PASS |
| Sterility test | TSB 20-25°C | Absence of growth After 3 months at -20°C | Absence of growth | PASS |
| Electrophoretical pattern | SDS-PAGE | 1 band between 12 and 16 kDa | 1 band between 12 and 16 kDa | PASS |
| Identity | Western Blot | 1 band between 12 and 16 kDa | 1 band between 12 and 16 kDa | PASS |
| Activity | Bioassay | > 5 10^6 U/mg | 11,8 10^6 U/mg | PASS |
| | | After 6 months at -20°C | _ | _ |
| Electrophoretical pattern | SDS-PAGE | 1 band between 12 and 16 kDa | 1 band between 12 and 15 kDa | PASS |
| Identity | Western Blot | 1 band between 12 and 16 kDa | 1 band between 12 and 15 kDa | PASS |
| Activity | Bioassay | > 5 10^6 U/mg | 12,7 10^6 U/mg | PASS |
| | | After 9 months at -20°C | | |
| Electrophoretical pattern | SDS-PAGE | 1 band between 12 and 16 kDa | 1 band between 12 and 15 kDa | PASS |

| Identity | Western Blot | 1 band between 12 and 16 kDa | 1 band between 12 and 15 kDa | PASS |
|---------------------------|----------------|------------------------------|------------------------------|------|
| Activity | Bioassay | > 5 10^6 U/mg | 12,0 10^6 U/mg | PASS |
| | | After 12 months at -20°C | | |
| Electrophoretical pattern | SDS-PAGE | 1 band between 12 and 16 kDa | 1 band between 12 and 15 kDa | PASS |
| Identity | Western Blot | 1 band between 12 and 16 kDa | 1 band between 12 and 15 kDa | PASS |
| Activity | Bioassay | > 5 10^6 U/mg | 13,8 10^6 U/mg | PASS |
| | | After 18 months at -20°C | | |
| Electrophoretical pattern | SDS-PAGE | 1 band between 12 and 16 kDa | 1 band between 12 and 15 kDa | PASS |
| Identity | Western Blot | 1 band between 12 and 16 kDa | 1 band between 12 and 15 kDa | PASS |
| Activity | Bioassay | > 5 10^6 U/mg | 13,2 10^6 U/mg | PASS |
| | | After 24 months at -20°C | | |
| Electrophoretical pattern | SDS-PAGE | 1 band between 12 and 16 kDa | 1 band between 12 and 15 kDa | PASS |
| Identity | Western Blot | 1 band between 12 and 16 kDa | 1 band between 12 and 15 kDa | PASS |
| Abnormal toxicity / | on mice | No weight loss, no abnormal | Absence | PASS |
| General safety | | Reaction | | |
| Abnormal toxicity / | on guinea-pigs | No weight loss, no abnormal | Absence | PASS |
| General safety | | Reaction | | |
| Sterility test | FTM 30-35°C | Absence of growth | Absence of growth | PASS |
| Sterility test | TSB 20-25°C | Absence of growth | Absence of growth | PASS |

GENTAUR rh IL-4 is manufactured in full compliance with cGMP in facilities approved by the Belgian Ministry of Health for the production and storage of medicinal products. GMP production at HENOGEN SA and GMP Lyophilisation at GSK Inc, Rixensart.



Use

GENTAUR rh IL-4 is not an approved medicinal product and cannot be injected as such to patients. However this GMP IL-4 is used in clinical tests for DC Therapy today in Belgium, France, Denmark, USA and Japan. CE CERTIFIED FOR EX VIVO CELL CULTURE AND DC THERAPY CLINICAL TESTS

Brussels, 04/02/2010 GNT#04GMPHUIL4-50UG

Gentaur Molecular products Avenue de l'Armée 68/B4 B1040 Brussel Tel: +32 16 58 90 45 Fax: +32 16 50 90 45